

Ateneo de Manila University

Archium Ateneo

Mathematics Faculty Publications

Mathematics Department

2015

Some Properties of the Exchange Operator with respect to Structured Matrices defined by Indefinite Scalar Product Spaces

Hanz Martin C. Cheng

Roden Jason David

Ateneo de Manila University

Follow this and additional works at: <https://archium.ateneo.edu/mathematics-faculty-pubs>



Part of the [Algebraic Geometry Commons](#), and the [Discrete Mathematics and Combinatorics Commons](#)

Recommended Citation

Cheng, H.M.C., David, R.J. (2015). Some properties of the exchange operator with respect to structured matrices defined by indefinite scalar product spaces. *Electronic Journal of Linear Algebra*, 30, Article number 21, 304-311.

This Article is brought to you for free and open access by the Mathematics Department at Archium Ateneo. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of Archium Ateneo. For more information, please contact oadrcw.ls@ateneo.edu.



**SOME PROPERTIES OF THE EXCHANGE OPERATOR WITH
 RESPECT TO STRUCTURED MATRICES DEFINED BY
 INDEFINITE SCALAR PRODUCT SPACES***

HANZ MARTIN C. CHENG[†] AND RODEN JASON DAVID[†]

Abstract. The properties of the exchange operator on some types of matrices are explored in this paper. In particular, the properties of $exc(A, p, q)$, where A is a given structured matrix of size $(p + q) \times (p + q)$ and $exc : M \times \mathbb{N} \times \mathbb{N} \rightarrow M$ is the exchange operator are studied. This paper is a generalization of one of the results in [N.J. Higham. *J*-orthogonal matrices: Properties and generation. *SIAM Review*, 45:504–519, 2003.].

Key words. Scalar product spaces, Structured matrices, Exchange operator.

AMS subject classifications. 15B10, 15B57, 46C20.

1. Introduction. In Mackey, Mackey and Tisseur [1] a *scalar product* $\langle x, y \rangle_M$ defined by a non-singular matrix M is given by

$$\langle x, y \rangle_M = \begin{cases} x^T M y & \text{for bilinear forms,} \\ x^* M y & \text{for sesquilinear forms.} \end{cases}$$

The *adjoint* of a matrix A with respect to the scalar product, denoted by $A^{\star M}$ is defined by the property $\langle Ax, y \rangle_M = \langle x, A^{\star M} y \rangle_M$. It can be shown that the matrix adjoint is given explicitly by

$$A^{\star M} = \begin{cases} M^{-1} A^T M & \text{for bilinear forms,} \\ M^{-1} A^* M & \text{for sesquilinear forms.} \end{cases}$$

For the above cases, $*$ is used to denote the conjugate transpose. An automorphism group \mathbb{G} , Lie algebra \mathbb{L} and Jordan algebra \mathbb{J} is associated with the scalar product. They are namely defined by:

$$\begin{aligned} \mathbb{G} &:= \{G \in \mathbb{K}^{n \times n} : \langle Gx, Gy \rangle_M = \langle x, y \rangle_M \ \forall x, y \in \mathbb{K}^n\} \\ &= \{G \in \mathbb{K}^{n \times n} : G^{\star M} = G^{-1}\}, \end{aligned}$$

*Received by the editors on February 8, 2015. Accepted for publication on May 8, 2015. Handling Editor: Michael Tsatsomeros.

[†]Department of Mathematics, Ateneo de Manila University, Quezon City, Philippines (hcheng@ateneo.edu, rj david@ateneo.edu).

$$\begin{aligned} \mathbb{L} &:= \{L \in \mathbb{K}^{n \times n} : \langle Lx, y \rangle_M = -\langle x, Ly \rangle_M \quad \forall x, y \in \mathbb{K}^n\} \\ &= \{L \in \mathbb{K}^{n \times n} : L^{\star M} = -L\}, \end{aligned}$$

$$\begin{aligned} \mathbb{J} &:= \{J \in \mathbb{K}^{n \times n} : \langle Jx, y \rangle_M = \langle x, Jy \rangle_M \quad \forall x, y \in \mathbb{K}^n\} \\ &= \{J \in \mathbb{K}^{n \times n} : J^{\star M} = J\}, \end{aligned}$$

where $\mathbb{K} = \mathbb{R}$ or \mathbb{C} .

For this work, the automorphism group, Jordan algebra, and Lie algebra with respect to a scalar product defined by the nonsingular matrix M will be denoted as \mathbb{G}_M , \mathbb{J}_M , and \mathbb{L}_M , respectively. Let n be a positive integer and let p, q be non-negative integers such that $p + q = n$. Let

$$\Sigma_{p,q} = \begin{bmatrix} I_p & O \\ O & -I_q \end{bmatrix}.$$

A matrix $Q \in \mathbb{R}^{n \times n}$ is said to be *J-orthogonal* if and only if $Q = \Sigma_{p,q} Q^T \Sigma_{p,q}$. A matrix $A \in \mathbb{R}^{n \times n}$ can be partitioned in the form

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix},$$

where $A_{11} \in \mathbb{R}^{p \times p}$, $A_{12} \in \mathbb{R}^{p \times q}$, $A_{21} \in \mathbb{R}^{q \times p}$, and $A_{22} \in \mathbb{R}^{q \times q}$. If A_{11} is nonsingular, then we define the *exchange operator* as

$$exc(A, p, q) = \begin{bmatrix} A_{11}^{-1} & -A_{11}^{-1} A_{12} \\ A_{21} A_{11}^{-1} & A_{22} - A_{21} A_{11}^{-1} A_{12} \end{bmatrix}.$$

In some contexts, $exc(A, p, q)$ is also known as the principal pivot transform of A relative to A_{11} . Let $B = exc(A, p, q)$. In Tsatsomeros [3] the *exchange operator* $exc(A, p, q)$ is related to A such that for every $x = (x_1^T, x_2^T)^T$ and $y = (y_1^T, y_2^T)^T$ in \mathbb{C}^n partitioned conformally to A ,

$$A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$

if and only if

$$B \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}.$$

2. Results.

THEOREM 2.1 (Higham [4]). *Let $A \in \mathbb{R}^{n \times n}$. If A is J-orthogonal then $exc(A, p, q)$ is orthogonal. If A is orthogonal and A_{11} is nonsingular, then $exc(A, p, q)$ is J-orthogonal.*

First, we note that every J -orthogonal matrix is an element of the automorphism group $\mathbb{G}_{\Sigma_{p,q}}$ and every orthogonal matrix is an element of the automorphism group \mathbb{G}_{I_n} . Thus, Higham's theorem is equivalent to stating that: if A is an element of the automorphism group $\mathbb{G}_{\Sigma_{p,q}}$, then $exc(A, p, q) \in \mathbb{G}_{I_n}$. Also, if $A \in \mathbb{G}_{I_n}$, then it follows that $exc(A, p, q) \in \mathbb{G}_{\Sigma_{p,q}}$. We generalize the result above onto the automorphism group \mathbb{G}_M and extend it to the Jordan algebra \mathbb{J}_M and the Lie algebra \mathbb{L}_M , where

$$M = \begin{bmatrix} N_1 & O \\ O & N_2 \end{bmatrix}$$

is a matrix such that $N_1 \in \mathbb{R}^{p \times p}$ and $N_2 \in \mathbb{R}^{q \times q}$ are both non-singular. Let

$$\hat{M} = \begin{bmatrix} N_1 & O \\ O & -N_2 \end{bmatrix}.$$

THEOREM 2.2. *If $A \in \mathbb{G}_M$, then $exc(A, p, q) \in \mathbb{G}_{\hat{M}}$. Also, if $A \in \mathbb{G}_{\hat{M}}$ then $exc(A, p, q) \in \mathbb{G}_M$.*

THEOREM 2.3. *If $A \in \mathbb{J}_M$, then $exc(A, p, q) \in \mathbb{J}_{\hat{M}}$. Also, if $A \in \mathbb{J}_{\hat{M}}$ then $exc(A, p, q) \in \mathbb{J}_M$.*

THEOREM 2.4. *The following statements:*

- a. $A \in \mathbb{L}_M \implies exc(A, p, q) \in \mathbb{L}_{\hat{M}}$,
- b. $A \in \mathbb{L}_{\hat{M}} \implies exc(A, p, q) \in \mathbb{L}_M$

are true if and only if A is of the form $\begin{bmatrix} A_{11} & O_{p \times q} \\ O_{q \times p} & A_{22} \end{bmatrix}$.

We only prove Theorem 2.2 since the proof of Theorem 2.3 and Theorem 2.4 can be done similarly.

Proof of Theorem 2.2. $A \in \mathbb{G}_M \rightarrow A^{\star M} = A^{-1} \text{ or } A^{\star M} A = I$. Taking the product, we get

$$(2.1) \quad N_1^{-1} A_{11}^* N_1 A_{11} + N_1^{-1} A_{21}^* N_2 A_{21} = I,$$

$$(2.2) \quad N_2^{-1} A_{12}^* N_1 A_{11} + N_2^{-1} A_{22}^* N_2 A_{21} = O,$$

$$(2.3) \quad N_1^{-1} A_{11}^* N_1 A_{12} + N_1^{-1} A_{21}^* N_2 A_{22} = O,$$

$$(2.4) \quad N_2^{-1} A_{12}^* N_1 A_{12} + N_2^{-1} A_{22}^* N_2 A_{22} = I.$$

We need to show that the product $exc(A, p, q)^{\star \hat{M}} exc(A, p, q) = I$.

The A_{11} block is given by

$$\begin{aligned} N_1^{-1} A_{11}^{-*} N_1 A_{11}^{-1} - N_1^{-1} A_{11}^{-*} A_{21}^* N_2 A_{21} A_{11}^{-1} \\ = N_1^{-1} A_{11}^{-*} N_1 A_{11}^{-1} - N_1^{-1} A_{11}^{-*} (N_1 A_{11}^{-1} - A_{11}^* N_1) \text{ by (2.1)} \\ = I. \end{aligned}$$

The A_{12} block is given by

$$\begin{aligned}
 & -N_1^{-1}A_{11}^{-*}N_1A_{11}^{-1}A_{12} - N_1^{-1}A_{11}^{-*}A_{21}^*N_2(A_{22} - A_{21}A_{11}^{-1}A_{12}) \\
 & = -N_1^{-1}A_{11}^{-*}N_1A_{11}^{-1}A_{12} - N_1^{-1}A_{11}^{-*}(-A_{11}^*N_1A_{12}) \\
 & \quad + N_1^{-1}A_{11}^{-*}A_{21}^*N_2A_{21}A_{11}^{-1}A_{12} \text{ by (2.3)} \\
 & = -N_1^{-1}A_{11}^{-*}N_1A_{11}^{-1}A_{12} + A_{12} \\
 & \quad + N_1^{-1}A_{11}^{-*}(N_1 - A_{11}^*N_1A_{11})A_{11}^{-1}A_{12} \\
 & = O.
 \end{aligned}$$

The A_{21} block is given by

$$\begin{aligned}
 & N_2^{-1}A_{12}^*A_{11}^{-*}N_1A_{11}^{-1} + N_2^{-1}A_{22}^*N_2A_{21}A_{11}^{-1} - N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2A_{21}A_{11}^{-1} \\
 & = N_2^{-1}A_{12}^*A_{11}^{-*}N_1A_{11}^{-1} + (-N_2^{-1}A_{12}^*N_1A_{11})A_{11}^{-1} \\
 & \quad - N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2A_{21}A_{11}^{-1} \text{ by (2.2)} \\
 & = N_2^{-1}A_{12}^*A_{11}^{-*}N_1A_{11}^{-1} - N_2^{-1}A_{12}^*N_1 \\
 & \quad - N_2^{-1}A_{12}^*A_{11}^{-*}(N_1 - A_{11}^*N_1A_{11})A_{11}^{-1} \text{ by (2.1)} \\
 & = O.
 \end{aligned}$$

Finally, the A_{22} block is given by

$$\begin{aligned}
 & -N_2^{-1}A_{12}^*A_{11}^{-*}N_1A_{11}^{-1}A_{12} + N_2^{-1}A_{22}^*N_2A_{22} - N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2A_{22} \\
 & \quad - N_2^{-1}A_{22}^*N_2A_{21}A_{11}^{-1}A_{12} + N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2A_{21}A_{11}^{-1}A_{12} \\
 & = -N_2^{-1}A_{12}^*A_{11}^{-*}N_1A_{11}^{-1}A_{12} + N_2^{-1}A_{22}^*N_2A_{22} \\
 & \quad - N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2A_{22} - N_2^{-1}A_{22}^*N_2A_{21}A_{11}^{-1}A_{12} \\
 & \quad + N_2^{-1}A_{12}^*A_{11}^{-*}(N_1 - A_{11}^*N_1A_{11})A_{11}^{-1}A_{12} \text{ by (2.1)} \\
 & = N_2^{-1}A_{22}^*N_2A_{22} - N_2^{-1}A_{12}^*A_{11}^{-*}(-A_{11}^*N_1A_{12}) \\
 & \quad - N_2^{-1}(-A_{12}^*N_1A_{11})A_{11}^{-1}A_{12} - N_2^{-1}A_{12}^*N_1A_{12} \\
 & \quad \text{by (2.2) and (2.3)} \\
 & = I \text{ by (2.4)}.
 \end{aligned}$$

Thus, $\text{exc}(A, p, q) \in \mathbb{G}_{\hat{M}}$. The converse can be proven in a similar manner. \square

Let

$$K = \begin{bmatrix} O & N_1 \\ N_2 & O \end{bmatrix}, \quad \hat{K} = \begin{bmatrix} O & N_1 \\ -N_2 & O \end{bmatrix},$$

where $N_1 \in \mathbb{R}^{p \times p}$ and $N_2 \in \mathbb{R}^{q \times q}$ are both non-singular.

THEOREM 2.5. *If $A \in \mathbb{G}_K$, then $\text{exc}(A, p, q) \in \mathbb{J}_{\hat{K}}$.*

THEOREM 2.6. *If $A \in \mathbb{J}_K$, then $\text{exc}(A, p, q) \in \mathbb{G}_{\hat{K}}$.*

We only prove Theorem 2.5 since the proof of Theorem 2.6 can be done similarly.

Proof of Theorem 2.5. $A \in \mathbb{G}_K \implies A^{\star\kappa} = A^{-1}$ or $A^{\star\kappa}A = I$. Taking the product, we get

$$(2.5) \quad N_2^{-1}A_{22}^*N_2A_{11} + N_2^{-1}A_{12}^*N_1A_{21} = I,$$

$$(2.6) \quad N_1^{-1}A_{21}^*N_2A_{11} + N_1^{-1}A_{11}^*N_1A_{21} = O,$$

$$(2.7) \quad N_2^{-1}A_{22}^*N_2A_{12} + N_2^{-1}A_{12}^*N_1A_{22} = O,$$

$$(2.8) \quad N_1^{-1}A_{21}^*N_2A_{12} + N_1^{-1}A_{11}^*N_1A_{22} = I.$$

We need to show that $\hat{K}^{-1}exc(A, p, q)^*\hat{K} = exc(A, p, q)$. Consider

$$\begin{aligned} & \begin{bmatrix} O & -N_2^{-1} \\ N_1^{-1} & O \end{bmatrix} \begin{bmatrix} A_{11}^{-*} & A_{11}^{-*}A_{21}^* \\ -A_{12}^*A_{11}^{-*} & A_{22}^* - A_{12}^*A_{11}^{-*}A_{21}^* \end{bmatrix} \begin{bmatrix} O & N_1 \\ -N_2 & O \end{bmatrix} \\ &= \begin{bmatrix} N_2^{-1}A_{12}^*A_{11}^{-*} & -N_2^{-1}A_{22}^* + N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^* \\ N_1^{-1}A_{11}^{-*} & N_1^{-1}A_{11}^{-*}A_{21}^* \end{bmatrix} \begin{bmatrix} O & N_1 \\ -N_2 & O \end{bmatrix} \\ &= \begin{bmatrix} N_2^{-1}A_{22}^*N_2 - N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2 & N_2^{-1}A_{12}^*A_{11}^{-*}N_1 \\ -N_1^{-1}A_{11}^{-*}A_{21}^*N_2 & N_1^{-1}A_{11}^{-*}N_1 \end{bmatrix}. \end{aligned}$$

Using equations (2.5)–(2.8), we have

$$\begin{aligned} & N_2^{-1}A_{22}^*N_2 - N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2 \\ &= A_{11}^{-1} - N_2^{-1}A_{12}^*N_1A_{21}A_{11}^{-1} - N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2 \\ &= A_{11}. \end{aligned}$$

Also,

$$\begin{aligned} N_2^{-1}A_{12}^*A_{11}^{-*}N_1 &= N_2^{-1}A_{12}^*(A_{11}^{-*}A_{21}^*N_2A_{12} + N_1A_{22}) \\ &= N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2A_{12} + N_2^{-1}A_{12}^*N_1A_{22} \\ &= N_2^{-1}A_{12}^*(A_{11}^{-*}A_{21}^*N_2)A_{12} - N_2^{-1}A_{22}^*N_2A_{12} \\ &= N_2^{-1}A_{12}^*(-N_1A_{21}A_{11}^{-1})A_{12} - (A_{11}^{-1} - N_2^{-1}A_{12}^*N_1A_{21}A_{11}^{-1})A_{12} \\ &= -A_{11}^{-1}A_{12} \end{aligned}$$

and

$$\begin{aligned} -N_1^{-1}A_{11}^{-*}A_{21}^*N_2 &= -N_1^{-1}(-N_1A_{21}A_{11}^{-1}) \\ &= A_{21}A_{11}^{-1}. \end{aligned}$$

Finally,

$$\begin{aligned} N_1^{-1}A_{11}^{-*}N_1 &= N_1^{-1}(N_1A_{22} + A_{11}^{-*}A_{21}^*N_2A_{12}) \\ &= A_{22} + N_1^{-1}(A_{11}^{-*}A_{21}^*N_2)A_{12} \\ &= A_{22} + N_1^{-1}(-N_1A_{21}A_{11}^{-1})A_{12} \\ &= A_{22} - A_{21}A_{11}^{-1}A_{12}. \end{aligned}$$

Thus,

$$\begin{aligned} & \begin{bmatrix} N_2^{-1}A_{22}^*N_2 - N_2^{-1}A_{12}^*A_{11}^{-*}A_{21}^*N_2 & N_2^{-1}A_{12}^*A_{11}^{-*}N_1 \\ -N_1^{-1}A_{11}^{-*}A_{21}^*N_2 & N_1^{-1}A_{11}^{-*}N_1 \end{bmatrix} \\ &= \begin{bmatrix} A_{11}^{-1} & -A_{11}^{-1}A_{12} \\ A_{21}A_{11}^{-1} & A_{22} - A_{21}A_{11}^{-1}A_{12} \end{bmatrix} \end{aligned}$$

or equivalently, $\hat{K}^{-1}exc(A, p, q)^*\hat{K} = exc(A, p, q)$. \square

3. Application (The Schur complement method). Matrices that can be directly factored are limited in size due to large memory requirements. Thus, the difficulty of solving large-scale linear systems of equations using standard techniques arises. Preconditioned iterative solvers require less memory, but often suffer from slow convergence. One of the ideas that have been developed to address these problems is called the Schur complement method. Several parallel hybrid solvers have been developed based on the above mentioned idea [2].

In the Schur complement method, instead of solving the system

$$(3.1) \quad \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$

directly, we consider

$$\begin{bmatrix} A_{11}^{-1} & -A_{11}^{-1}A_{12} \\ A_{21}A_{11}^{-1} & A_{22} - A_{21}A_{11}^{-1}A_{12} \end{bmatrix} \begin{bmatrix} y_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \\ y_2 \end{bmatrix}$$

and solve the system of equations

$$\begin{aligned} A_{11}^{-1}y_1 - A_{11}^{-1}A_{12}x_2 &= x_1, \\ A_{21}A_{11}^{-1}y_1 + (A_{22} - A_{21}A_{11}^{-1}A_{12})x_2 &= y_2. \end{aligned}$$

In solving the above system, we need to compute $(A_{22} - A_{21}A_{11}^{-1}A_{12})^{-1}$.

Suppose that p and q are integers such that $q \leq p$ and

$$M = \begin{bmatrix} N_1 & O \\ O & N_2 \end{bmatrix}$$

is a matrix such that $N_1 \in \mathbb{R}^{p \times p}$ and $N_2 \in \mathbb{R}^{q \times q}$ are both non-singular. Let

$$\hat{M} = \begin{bmatrix} N_1 & O \\ O & -N_2 \end{bmatrix}.$$

Suppose $A \in \mathbb{G}_M$. Since $exc(A, p, q) \in \mathbb{G}_{\hat{M}}$ by Theorem 2.2, we know that $(A_{22} - A_{21}A_{11}^{-1}A_{12})N_2^{-*}(-A_{11}^{-1}A_{12}) = A_{21}A_{11}^{-1}N_1^{-*}A_{11}^{-*}$. Suppose further that $rank(A_{21}) =$

q. Then we may compute

$$\begin{aligned} & (A_{22} - A_{21}A_{11}^{-1}A_{12})^{-1} \\ &= N_2^{-*}(-A_{11}^{-1}A_{12})^*(A_{11}^{-1})^{-*}N_1^*(A_{21}A_{11}^{-1})^*((A_{21}A_{11}^{-1})(A_{21}A_{11}^{-1})^*)^{-1} \\ &= -N_2^{-*}A_{12}^*N_1^*A_{11}^{-*}A_{21}^*(A_{21}A_{11}^{-1}A_{11}^{-*}A_{21}^*)^{-1} \end{aligned}$$

explicitly.

To compute $(A_{21}A_{11}^{-1}A_{11}^{-*}A_{21}^*)^{-1}$, we perform singular value decomposition on both A_{11} and A_{21} . Write $A_{11} = U\Sigma V^*$ and $A_{21} = \hat{U}\hat{\Sigma}\hat{V}^*$, where U, V, \hat{U} , and \hat{V} are unitary and

$$\Sigma = \begin{bmatrix} \sigma_1 & 0 & \dots & 0 \\ 0 & \sigma_2 & \dots & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \sigma_p \end{bmatrix}, \quad \hat{\Sigma} = \begin{bmatrix} \hat{\sigma}_1 & 0 & 0 & \dots & 0 & 0 \\ 0 & \hat{\sigma}_2 & 0 & \dots & 0 & 0 \\ 0 & 0 & \ddots & \dots & 0 & 0 \\ 0 & 0 & 0 & \hat{\sigma}_q & \dots & 0 \end{bmatrix}$$

such that $\{\sigma_1, \sigma_2, \dots, \sigma_p\}$ and $\{\hat{\sigma}_1, \hat{\sigma}_2, \dots, \hat{\sigma}_q\}$ are the singular values of A_{11} and A_{21} , respectively. Then,

$$\begin{aligned} A_{21}A_{11}^{-1}A_{11}^{-*}A_{21}^* &= \hat{U}\hat{\Sigma}\hat{V}^*V\Sigma^{-1}U^*U\Sigma^{-*}V^*\hat{V}\hat{\Sigma}^*\hat{U}^* \\ &= \hat{U}\hat{\Sigma}\hat{V}^*V\Sigma^{-1}\Sigma^{-*}V^*\hat{V}\hat{\Sigma}^*\hat{U}^*. \end{aligned}$$

Finally,

$$\begin{aligned} (A_{21}A_{11}^{-1}A_{11}^{-*}A_{21}^*)^{-1} &= \hat{U}(\hat{\Sigma}^+)^*\hat{V}^*V\Sigma^*\Sigma V^*\hat{V}\hat{\Sigma}^+\hat{U}^* \\ &= (\Sigma V^*\hat{V}\hat{\Sigma}^+\hat{U}^*)^*(\Sigma V^*\hat{V}\hat{\Sigma}^+\hat{U}^*), \end{aligned}$$

where

$$\hat{\Sigma}^+ = \begin{bmatrix} \frac{1}{\hat{\sigma}_1} & 0 & \dots & 0 \\ 0 & \frac{1}{\hat{\sigma}_2} & \dots & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \frac{1}{\hat{\sigma}_q} \\ 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Thus, given $A \in \mathbb{G}_M$, the solution of (3.1) is given explicitly by

$$\begin{aligned} x_2 &= (\Sigma V^*\hat{V}\hat{\Sigma}^+\hat{U}^*)^*(\Sigma V^*\hat{V}\hat{\Sigma}^+\hat{U}^*)(y_2 - A_{21}A_{11}^{-1}y_1), \\ x_1 &= A_{11}^{-1}y_1 - A_{11}^{-1}A_{12}x_2. \end{aligned}$$

REFERENCES

- [1] D.S. Mackey, N. Mackey, and F. Tisseur. On the definition of two natural classes of scalar product. *MIMS EPrints* 2007.64, Manchester Institute for Mathematical Sciences, The University of Manchester, 2007.
- [2] I. Yamazaki and X.S. Li. On techniques to improve robustness and scalability of the Schur complement method. In: J.M. Laginha, M. Palma, M. Daydé, O. Marques, and J. Correia Lopes (editors), *High Performance Computing for Computational Science - VECPAR 2010*, Springer-Verlag, Berlin-Heidelberg, 421–434, 2011.
- [3] M.J. Tsatsomeros. Principal pivot transforms: Properties and applications. *Linear Algebra and its Applications*, 307:151–165, 2000.
- [4] N.J. Higham. J -orthogonal matrices: Properties and generation. *SIAM Review*, 45:504–519, 2003.