Ateneo de Manila University

### Archium Ateneo

**Biology Faculty Publications** 

**Biology Department** 

4-2014

# Effects of replacing fishmeal with squash seed meal (Cucurbita maxima) on performance of juvenile Nile tilapia (Oreochromis niloticus)

Janice A. Ragaza Ateneo de Manila University

Jonni Fay C. Teves

Marie Noelle Fernandez Osaka University

Follow this and additional works at: https://archium.ateneo.edu/biology-faculty-pubs

Part of the Aquaculture and Fisheries Commons, and the Biology Commons

### **Custom Citation**

Ragaza, Janice & Teves, Jonni & Fernandez, Marie Noelle. (2014). Effects of replacing fishmeal with squash seed meal (Cucurbita maxima) on performance of juvenile Nile tilapia (Oreochromis niloticus). AACL Bioflux. 7. 68-75.

This Article is brought to you for free and open access by the Biology Department at Archīum Ateneo. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of Archīum Ateneo. For more information, please contact oadrcw.ls@ateneo.edu.

# AACL BIOFLUX

### Aquaculture, Aquarium, Conservation & Legislation International Journal of the Bioflux Society

## Effects of replacing fishmeal with squash seed meal (*Cucurbita maxima*) on performance of juvenile Nile tilapia (*Oreochromis niloticus*)

Jonni Fay C. Teves, Marie Noelle M. Fernandez, Janice A. Ragaza

Department of Biology, School of Science and Engineering, Loyola Schools, Ateneo de Manila University, Katipunan Avenue, Loyola Heights, Quezon City 1108, Metro Manila, Philippines. Corresponding author: J. A. Ragaza, jragaza@ateneo.edu

**Abstract**. The effects of replacing fishmeal with squash (*Cucurbita maxima*) seed meal (SSM) on *Oreochromis niloticus* juveniles were investigated on a 30-day feeding trial. Triplicate groups of ten (10) *O. niloticus* juveniles ( $2.54\pm0.949$  g) stocked in polyethylene tanks received iso-nitrogenous diets with SSM inclusion levels of 0%, 5%, 10%, 15% and 20%. Fish fed 5% SSM inclusion level showed the highest average weight gain ( $202.25\pm36.26\%$ ) and specific growth rate ( $3.76\pm0.42\%$ ) among the fish fed SSM-containing diets. However, there were no significant differences between the average weight gain and specific growth rate of fish in all diets. Fish fed S5, S15 and S20 showed significantly higher protein carcass levels; lipid levels were not significantly different while hepatosomatic index in fish fed S20 was significantly higher than fish fed other diets. Cost analysis of diets revealed S5 as the most cost-effective diet. SSM diets were comparable to the positive control diet as these were efficiently utilized by the fish, thereby making it a potential candidate ingredient/replacement for fishmeal. **Key Words**: *Cucurbita maxima* seeds, *Oreochromis niloticus*, growth performance, feed utilization efficiency, carcass composition.

**Introduction**. The aquaculture sector is vital to maintain fish supply, especially with the decline in capture fish industry due to unsustainable practices (FAO 2001; FAO 2006). However, aquaculture, which is highly dependent on fishmeal as a protein source for fish feeds, also proves to be no longer sustainable (Tacon & Metian 2008). The steady supply of fishmeal and fish oil vis-à-vis the rapid expansion of aquaculture has contributed to a rise in costs of production, and in effect a rise in the cost of fish. This might be one of the factors associated with the continuous rise in the retail price of the low-priced fish Oreochromis niloticus (Nile tilapia), one of the most dominant, farmed aquaculture species in the Philippines, and is also the subject of this research (EI-Sayed 2006; BAS 2013; Bahnasawy 2009). The rise of aquaculture production is thus highly important to fill in the gap between supply and demand, so as to keep the price of food fish at a reasonable range for the rural and urban poor which is critical for food security (Huntington & Hasan 2009). In this context, the formulation of fish feed using a cheap, locally available and highly nutritious terrestrial resource such as Cucurbita maxima seeds is needed. Crude protein levels of C. maxima seeds and kernels are comparable to high protein-containing seeds and legumes such as soybeans and cowpea; while lipid content of C. maxima kernels is comparable to sunflower, soybeans and cotton seeds (Alfawaz 2004; Hayden 1990; Srbinoska et al 2012; Haciseferogullari & Acaroglu 2012). Seed oils are mainly composed of polyunsaturated fatty acids, namely oleic and linoleic acids (Radovich et al 2010; Kirbaslar et al 2012) with the latter accounting for more than half of the seed's fatty acid content (Kim et al 2012). Its seeds also contain vitamins A, C and E (Raganathan et al 2013), calcium, iron, phosphorous and zinc, and cucurbitacins (Khare 2007; Glew et al 2006). Amino acids detected in significant amounts were glutamic acid, aspartic acid, leucine, glycine, valine, among others (Kim et al 2012).

To our knowledge, there is no reported study on the inclusion of squash seeds in fish feed. However, formulation of *C. moschata* seed meal, which belongs to the same

plant family, has been made for broilers and has shown to reduce serum triglyceride and cholesterol levels due to its high oleic acid content compared to conventional seeds (Aguilar et al 2011). Squash leaves which contain significant amount of protein have also been tested as plant protein source for O. niloticus fingerlings diets. It resulted to lowest weight gain compared to control diets although within the acceptable survival rate of 98% (Magouz et al 2008).

The high nutrition content of C. maxima seeds can be a potential replacement to high-protein fishmeal, thereby reduce reliance on marine fish species and help in its recovery from overexploitation and depletion. Rising costs associated with fish feed production can also be alleviated, which can make cheaper, protein-rich fish staples available to a greater number of people.

This study was conducted to determine the effects of different partial replacement levels of squash seed meal (SSM) on the growth performance, feed utilization efficiency and carcass composition of O. niloticus juveniles.

Material and Method. A feeding trial was conducted in the months of July to August 2013. In preparation for the feeding trial, C. maxima seeds authenticated by Dr. James V. LaFankie of the Institute of Biology Herbarium, University of the Philippines, Diliman, Quezon City were washed with tap water, dried for a total of 30 minutes: 20 minutes at 150°C and subsequently 10 minutes at 125°C; air-dried for 48 hours and milled to 1 mm powder. Iso-nitrogenous diets with SSM inclusion levels of 0%, 5%, 10% 15% and 20% were prepared at the Southeast Asian Fisheries Development Center (SEAFDEC), Binangonan Freshwater Station, Rizal. Diet with 0% SSM inclusion (S0) served as positive control. The following ingredients were used: local sardine fishmeal, shrimp meal and SSM as protein sources; soy lecithin and cod liver oil as lipid sources; and carboxymethyl cellulose (CMC) and cornstarch as carbohydrate, binder and filler sources (Table 1). Diets were pelletized into 1.5 mm in diameter and contained protein levels 41.15-41.95%, lipid levels 11.4-11.88%, and carbohydrate levels 24.24-24.67%. Dried pellets were stored at -4°C in a refrigerator until used.

Table 1

| Diet ingredient (%)                     | SO    | S5    | S10   | S15   | S20   |
|---|-------|-------|-------|-------|-------|
| Sardine fishmeal <sup>*</sup>           | 52    | 41.25 | 36.25 | 30.5  | 27    |
| SSM <sup>₹</sup>                        | 0     | 7.25  | 13    | 19    | 22    |
| Shrimp meal <sup>ŦŦ</sup>               | 11.25 | 18.5  | 20.75 | 23    | 24.5  |
| Cod liver oil <sup>¶</sup>              | 10.75 | 7     | 4     | 1.5   | 0.5   |
| Soy lecithin <sup>¶¶</sup>              | 1     | 1     | 1     | 1     | 1     |
| Cornstarch                              | 16    | 16    | 16    | 16    | 16    |
| Vitamin-mineral premix <sup>ø</sup>     | 6     | 6     | 6     | 6     | 6     |
| Vitamin C                               | 0.1   | 0.1   | 0.1   | 0.1   | 0.1   |
| CMC                                     | 2.9   | 2.9   | 2.9   | 2.9   | 2.9   |
| TOTAL                                   | 100   | 100   | 100   | 100   | 100   |
| Proximate composition (in % dry matter) |       |       |       |       |       |
| Crude protein                           | 41.86 | 41.95 | 41.95 | 41.52 | 41.15 |
| Crude lipid                             | 11.88 | 11.88 | 11.40 | 11.52 | 11.84 |
| Carbohydrate                            | 24.25 | 24.67 | 24.61 | 24.49 | 24.34 |

Formulation and proximate composition of test diets for partial replacement of fishmeal with SSM

\*SEAFDEC, Philippines; protein, 65.65; lipid, 0.82; carbohydrate, 15.86; ash, 16.97;

<sup>†</sup>Farmer's Market, Quezon City, Philippines; protein, 30.10; lipid, 43; carbohydrate, 10 (as reported by Alfawaz 2004; Srbinoska et al 2012; Kim et al 2012); <sup>++</sup>SEAFDEC, Philippines; protein, 68.6; lipid, 3.9; carbohydrate, 7.6;

<sup>¶</sup>SEAFDEC, Philippines; lipid, 100;

<sup>¶¶</sup>SEAFDEC, Philippines; lipid, 70;

<sup>o</sup>Cual, Kuan and Cual International Trading Corp., Quezon City. Mixture contains vitamins (A, B1, B2, B6, D3, E), folic acid, niacin, pathothenic acid, biotin, choline chloride, iron, copper, iodine, manganese, zinc, cobalt, and selenium.

**Experimental design**. A total of two hundred and twenty-five (225) *O. niloticus* juveniles were acclimatized for 1 week in polyethylene tanks and fed on commercial diet Neon Tetra Guppy (Sealion NTG-60) with a minimum of 43% crude protein, minimum of 3% crude fat, maximum of 6% crude fiber and maximum of 10% moisture to apparent satiation. The 30-day feeding trial ensued as  $2.54 \pm 0.949$  g (mean  $\pm$  standard error of the mean SEM) pre-weighed fingerlings were randomly distributed to fifteen 30-L capacity polyethylene tanks with a stocking density of 10 fingerlings per tank. Diets were randomly distributed among the fifteen (15) tanks with three replicates per treatment. Daily ration was divided equally into two meals at 08:00 h and 16:00 h, hand-fed to apparent satiation. Tanks were installed with individual filtration systems. Removal of wastes and replenishment of 40% rearing water were done daily to maintain optimum water quality. Uneaten feeds were removed, freeze-dried and weighed to calculate for corrected feed intake. Sampling was conducted once every two weeks to monitor growth performance.

**Growth performance and feed utilization**. At the end of the 30-day feeding trial, growth parameters of *O. niloticus* were evaluated and expressed as percent average weight gain (% AWG), specific growth rate (SGR), feed conversion ratio (FCR), feed intake (FI) and protein efficiency ratio (PER).

**Sample collection and body composition of fish**. An initial sample of twenty-five (25) *O. niloticus* fingerlings from stock were individually weighed for biochemical composition analysis. Prior to final sampling, all fish were starved for 24 hours to eliminate traces of other prey or feeds in the digestive tract (Rowe & Dean 1998). Remaining fish per tank were collected from each treatment, individually weighed, anaesthetized by hypothermia treatment, freeze-dried and kept at -80°C until analyzed for whole body protein, crude lipid and moisture composition. After weighing, liver was removed, weighed and pooled to calculate for the hepatosomatic index (HSI).

**Proximate chemical analysis of fish and diets**. All analyses were done in triplicate. The proximate composition of fish carcass and diets were analyzed according to the standard methods of AOAC (1990). Moisture content was estimated by drying fish carcass to a constant weight at 105°C for 7 hours in a vacuum oven. These were then homogenized in a blender. Lipid content of 2 g of sample was analyzed via diethyl ether extraction using a Soxhlet apparatus for 4 h. Remaining fish carcass samples were sent to the Philippine Institute for Pure and Applied Chemistry (PIPAC), Loyola Heights, Quezon City for protein content analysis.

*Cost analysis of diets*. Parameters of cost analysis computed for included feed cost kg<sup>-1</sup> of diet, incidence cost and profit index.

**Statistical analysis**. Data were subjected to one-way analysis of variance (ANOVA) to test the effect of the diet as a factor and to see the difference between the groups. Tukey post-hoc test was conducted to compare between means at  $p \le 0.05$ . The IBM-SPSS v. 21 (IBM Corp., New York, USA) software was used for statistical analysis.

**Results and Discussion**. Fish fed 5% SSM inclusion level showed the highest % AWG (202.25  $\pm$  36.26%) and SGR (3.76  $\pm$  0.42%) among fish fed SSM-containing diets (Table 2). There were no significant differences between the % AWG and SGR of fish in all diets. Nonetheless, SSM inclusion in diets resulted in positive growth response in fish. FI decreased with increasing SSM inclusion, reaching 0.229  $\pm$  0.050 g fish<sup>-1</sup> day<sup>-1</sup> for fish fed S10, but increased to 0.268  $\pm$  0.051 g fish<sup>-1</sup> day<sup>-1</sup> for fish fed S20. This indicated that all diets, including those with SSM inclusion, were readily accepted and eaten by *O. niloticus* fingerlings. There was no feed rejection observed in all dietary treatments. FCR was found to be numerically lowest in fish fed S5 (0.048  $\pm$  0.013). Lower FCR denotes more efficient feed utilization, i.e., there is faster conversion of feed into weight gain of fish. This is economically favorable in fish farming. Even with low quantity of feeds, fish

growth performance is optimized, hence reducing production costs associated with fish feeding. Fish fed 5% SSM also showed numerically highest PER (0.181  $\pm$  0.04). High PER shows that smaller amount of protein intake is needed to induce weight gain in fish, and hence is indicative of protein quality of feeds. There were no significant differences between the FCR and PER of fish in all treatments.

Table 2

Growth performance and feed utilization efficiency of juvenile *O. niloticus* fed diets partially replaced with different levels of SSM for 30 days<sup>\*</sup>

| Diet | AWG<br>(%)         | SGR<br>(% day <sup>-1</sup> ) <sup>¶</sup> | FCR <sup>T</sup>  | FI<br>(g fish <sup>-1</sup> day <sup>-1</sup> ) | $PER^{\Psi}$     |
|------|--------------------|--|-------------------|---|------------------|
| S0   | 213.50 ± 69.49     | $3.78 \pm 0.74$                            | $0.056 \pm 0.013$ | $0.243 \pm 0.024$                               | $0.157 \pm 0.03$ |
| S5   | $202.25 \pm 36.26$ | $3.76 \pm 0.42$                            | $0.048 \pm 0.013$ | $0.234 \pm 0.036$                               | $0.181 \pm 0.04$ |
| S10  | 149.21 ± 28.67     | $3.10 \pm 0.41$                            | $0.061 \pm 0.006$ | $0.229 \pm 0.050$                               | $0.133 \pm 0.02$ |
| S15  | $114.64 \pm 30.30$ | $2.56 \pm 0.49$                            | $0.064 \pm 0.014$ | $0.239 \pm 0.006$                               | $0.103 \pm 0.03$ |
| S20  | 175.72 ± 18.22     | $3.47~\pm~0.29$                            | $0.057 \pm 0.004$ | 0.268 ± 0.051                                   | $0.142 \pm 0.01$ |

\*Each value is the mean ± SEM of data from triplicate groups. Within a column, absence of letters indicates no significant differences between treatments;

<sup>¶</sup>Specific growth rate = [100 x (In final fish weight - In initial fish weight)]/days;

<sup>T</sup>Feed conversion ratio = dry feed intake (g)/live weight gain (g);

<sup> $\Psi$ </sup>Protein efficiency ratio = live weight gain (g)/protein intake (g).

Carcass composition analysis was conducted to elucidate the effects of SSM replacement on utilization, digestibility and deposition in body tissues of nutrients by *O. niloticus* (Table 3).

Table 3

| Diet    | Moisture         | Protein              | Lipid            | HSI <sup>¶</sup>              |
|---------|------------------|----------------------|------------------|-------------------------------|
| Initial | 79.47            | $58.90 \pm 0.25$     | 15.60            | NDΨ                           |
| S0      | $73.21 \pm 0.11$ | $56.33 \pm 0.15^{a}$ | $20.26 \pm 2.16$ | $0.0252\ \pm\ 0.0086^a$       |
| S5      | $73.52 \pm 0.49$ | 57.50 <sup>b</sup>   | $18.50 \pm 2.24$ | $0.0341 \pm 0.0090a^{b}$      |
| S10     | $73.60 \pm 0.72$ | $55.93 \pm 0.37^{a}$ | $19.50 \pm 1.20$ | $0.0323 \pm 0.0091a^{b}$      |
| S15     | 75.93 ± 1.06     | 58.50 <sup>b</sup>   | $18.90 \pm 0.20$ | 0.0279 ± 0.0098a <sup>b</sup> |
| S20     | 72.29 ± 1.98     | $58.13 \pm 0.27^{b}$ | $22.77 \pm 1.71$ | $0.0351 \pm 0.0081^{b}$       |

<sup>\*</sup>Each value is the mean  $\pm$  SEM of data from triplicate groups. Within a column, means with the same letters are not statistically different (p > 0.05). Absence of letters indicates no significant difference between treatments. Initial values were not included in the statistical analysis;

<sup>¶</sup>HSI = liver weight x 100/body weight (Agbo 2008);

 $\Psi$ ND = Not determined.

Carcass protein levels ranged from 55.93-58.50%. SSM inclusion levels of S5, S15 and S20 showed significantly higher carcass protein levels compared to the S10 diet. It has been reported that plant protein-based diets can lead to lower nitrogen retention in salmonids as these diets contain lower digestible energy and suboptimal amino acid profile critical for muscle growth (Cheng et al 2003). This was not the case in the study, which could be explained by amino acid levels as the determining factor in meeting the metabolic demands of fish (Ayadi et al 2012). Aside from sufficient vitamin-mineral supplementation, comparable growth performance of fish fed SSM diets and FM-based diet could be attributed to the amino acid profile of *C. maxima* as reported by Kim et al (2012), which contributed to the overall nutritional profile of SSM in meeting the metabolic demands of *O. niloticus* juveniles. Furthermore, effect of fatty acid level supplementation on protein carcass level was investigated. It was presumed that diets with lower SSM inclusion contained higher linolenic acid (LNA) content as this mainly

comes from fish oils (cod liver oil) than linoleic acid (LA) from plant oils in higher SSM inclusion (Table 1). Omega-3 polyunsaturated fatty acids (LNA) can increase muscle anabolic signaling and protein synthesis (Smith et al 2011). Contrary to the results in this study, fish fed S0, which was expected to show highest protein level, actually showed significantly lower protein content than in fish fed other diets.

Fish fed S20 showed the numerically highest lipid level, although there were no significant differences in lipid levels among diets. It has been reported that replacement of fishmeal with plant protein sources could increase lipogenic enzyme activities in sea bass (*Dicentrarchus labrax*), which could lead to increased lipid and gross energy content in the whole body of fish at constant level of protein (Kaushik et al 2004). Overall, high nutrient deposition in fish fed higher SSM-inclusion diet levels could be a result of high feed intake, nutrient utilization and nutrient digestibility (Abdel-Tawwab et al 2010).

The hepatosomatic index describes the status of energy reserve in an animal (Sadekarpawar & Parikh 2013). Fish fed S0 showed numerically lowest HSI compared to fish fed SSM diets. Meanwhile, fish fed S20 showed significantly higher HSI than fish fed S0. Differences in HSI values are also a function of protein, lipid and carbohydrate levels in fish (Ali et al 2000). Significantly higher HSI in fish fed S20 could be attributed to the numerically highest lipid content in fish carcass, which may have contributed to a greater lipid accumulation in the liver (Yildiz 2004).

Cost analysis of diets was also conducted using the following parameters: feed cost, incidence cost and profit index. Input values to feed cost computation were individual feed ingredient costs, direct labor costs (oven drying, milling) and overhead costs (electricity, miscellaneous costs) (Table 4).

Table 4

| Ingredients                    | Unit cost kg <sup>-1</sup> (\$) | S0   | S5      | S10     | S15     | S20     |
|--------------------------------|---------------------------------|--|---------|---------|---------|---------|
|                                |                                 | Cost kg <sup>-1</sup> for each ingredient <sup>**</sup> (\$) |         |         |         |         |
| Sardine fishmeal               | 0.90                            | 0.47   | 0.37    | 0.32    | 0.27    | 0.24    |
| SSM                            | 0.62                            | 0  | 0.04    | 0.08    | 0.12    | 0.14    |
| Shrimp meal                    | 4.48                            | 0.5  | 0.83    | 0.93    | 1.03    | 1.1     |
| Cod liver oil                  | 6.83                            | 0.73   | 0.48    | 0.27    | 0.1     | 0.03    |
| Soy lecithin                   | 1.79                            | 0.02   | 0.02    | 0.02    | 0.02    | 0.02    |
| Cornstarch                     | 0.69                            | 0.11   | 0.11    | 0.11    | 0.11    | 0.11    |
| Vitamin-mineral premix         | 4.48                            | 0.27   | 0.27    | 0.27    | 0.27    | 0.27    |
| Vitamin C                      | 26.89                           | 0.03   | 0.03    | 0.03    | 0.03    | 0.03    |
| CMC                            | 2.49                            | 0.07   | 0.07    | 0.07    | 0.07    | 0.07    |
| Subtotal cost kg <sup>-1</sup> |                                 | 2.20   | 2.22    | 2.11    | 2.02    | 2.01    |
| Cost ton <sup>-1</sup>         |                                 | 2201.69  | 2218.84 | 2105.44 | 2021.04 | 2007.12 |
| Direct labor <sup>¶</sup>      |                                 | 19.22  | 19.22   | 19.22   | 19.22   | 19.22   |
|                                |                                 | Overhead costs   |         |         |         |         |
| Electricity§                   |                                 | 40.78  | 40.78   | 40.78   | 40.78   | 40.78   |
| $Miscellaneous^{\psi}$         |                                 | 111.05   | 111.90  | 106.23  | 102.01  | 101.32  |
| Total cost ton <sup>-1</sup>   |                                 | 2372.73  | 2390.74 | 2271.68 | 2183.05 | 2168.44 |
| Total cost kg <sup>-1</sup>    |                                 | 2.37   | 2.39    | 2.27    | 2.18    | 2.17    |

Computations of feed cost per diet\*

\*Based on method of computation by Agbayani (2002);

<sup>\*\*</sup>Cost kg<sup>-1</sup> for each of ingredient = (unit cost kg<sup>-1</sup>) (feed formulation composition in kg);

<sup>1</sup>Direct Labor = (minimum wage rate) (2 aides); where minimum wage rate for the agriculture sector in the Philippines as of 2013 = Php 429 or \$9.61;

<sup>§</sup>Electricity cost = (200 kWh) (\$0.204); where \$0.204 = price/kWh (Php 9.10);

<sup>m</sup>Miscellaneous cost = 5% (feed cost ton<sup>-1</sup> + direct labor cost);

Note: prices were calculated based on exchange rate value of 1 USD = Php 44.6324 as of March 3, 2014.

Price list of feed ingredients was provided by SEAFDEC, while SSM feed ingredient cost was calculated by adding heating cost (HC) and milling cost (MC) upon seed processing, with corresponding computations as follows:

HC kg<sup>-1</sup> = (wattage of convection oven)(total heating time)/dry weight of seeds in kg) (Price/kWh) =  $(1.14 \text{ kW}) (13.5h/5.408 \text{ kg}) (\$0.204) = \$0.58 \text{ kg}^{-1}$ ; MC kg<sup>-1</sup> = (wattage of miller)(total milling time) (Price/kWh) = (3.72 kW)(0.05h) $(\$0.204) = \$0.038 \text{ kg}^{-1};$ 

SSM cost  $kg^{-1} = HC + MC =$ \$0.62 kg<sup>-1</sup>.

Cost analysis revealed lowest incidence cost and highest profit index in the S5 diet (Table 5). A low incidence cost value means lower cost to produce a unit weight gain in fish. Meanwhile, a high profit index indicates a high amount of profit for every cost incurred in feeding. Altogether, these are important factors a fish farmer considers when choosing feed ingredients to minimize cost and maximize profit.

Table 5

|      | Cost analysis of diets (ba      | used on a 30-day period co       | omputation)               |
|------|---------------------------------|----------------------------------|---------------------------|
| Diet | Feed cost kg <sup>-1</sup> (\$) | Incidence cost (\$) <sup>¶</sup> | Profit index <sup>4</sup> |

| Diet | Feed cost kg <sup>-</sup> ' (\$) | Incidence cost (\$)" | Profit index <sup><math>\Phi</math></sup> |
|------|----------------------------------|----------------------|---|
| SO   | 2.37                             | 3.83                 | 0.57                                      |
| S5   | 2.39                             | 3.36                 | 0.66                                      |
| S10  | 2.27                             | 3.77                 | 0.58                                      |
| S15  | 2.18                             | 4.92                 | 0.45                                      |
| S20  | 2.17                             | 4.02                 | 0.55                                      |

<sup>¶</sup>IC = total cost of feed used/weight of the fish produced (Agbavani 2002); where total cost of feed = feed cost

 $kg^{-1}$  x feed intake;  $\Psi$ PI = value of fish/cost of feeding (Mensah & Attipoe 2013) using Tilapia retail price of Php 98.17 kg<sup>-1</sup> (BAS 2013) or \$2.20 kg<sup>-1</sup>.

Conclusions. Fish fed 5% SSM exhibited numerically highest growth performance and feed utilization efficiency. Moreover, diets with SSM inclusions of 5 to 20% were comparable to the positive control diet, as these were found to be efficiently utilized by O. niloticus fingerlings as well. Therefore, SSM may pose as a potential candidate ingredient for fishmeal replacement in O. niloticus feeds. For further study, it is recommended that the experimental period be extended to monitor more precisely the growth performance as well as behavioral interactions among fish that may affect their response to the diets. When more efficient means of defatting seeds is available, higher inclusion levels of SSM may be investigated to see whether or not it can yield better growth performance in fish. Further research on the effects of SSM on amino acid profiles of both diets and fish carcass is recommended. Moreover, squash kernels may also be tapped as these contain higher amounts of protein, and hence a good candidate for fishmeal replacement.

Acknowledgements. The authors are grateful to Dr. Frolan Aya of the Southeast Asian Fisheries Development Center (SEAFDEC), Binangonan Freshwater Station, Rizal for sharing his expertise during feed formulation and preparation. Mr. Edwin Palenzuela of the Ateneo Department of Biology is also acknowledged for his assistance on technical matters concerning the aquaculture laboratory; the faculty of the Ateneo Department of Biology for their invaluable comments and suggestions to improve this research paper; and Mr. Lance Go of the Ateneo Department of Chemistry for his supervision on chemical analysis procedures. This research was partially funded by the Sanggunian ng Mga Magaaral ng mga Paaralang Loyola ng Ateneo de Manila under the Sanggunian Subsidy System Program.

#### References

- Abdel-Tawwab M., Ahmad M. H., Khattab Y. A. E., Shalaby A. M. E., 2010 Effect of dietary protein level, initial body weight, and their interaction on the growth, feed utilization, and physiological alterations of Nile tilapia, *Oreochromis niloticus* (L.). Aquaculture 298:267-274.
- Agbayani R. F., 2002 Economics of feeding. In: Nutrition in tropical aquaculture: essentials of fish nutrition, feeds, and feeding of tropical aquatic species. Millamena O. M., Coloso R. M., Pascual F. P. (eds), SEAFDEC, Iloilo Philippines, pp. 209-217.
- Agbo N. W., 2008 Oilseed meals as dietary protein sources for juvenile Nile tilapia (*Oreochromis niloticus* L.) Thesis dissertiation. University of Stirling, 210 pp.
- Aguilar Y. M., Yero O. M., Navarro M. I. V., Hurtado C. A. B., Lopez J. A. C., Mejia L. B. G., 2011 Effect of squash seed meal (*Cucurbita moschata*) on broiler performance, sensory meat quality, and blood lipid profile. Brazilian Journal of Poultry Science 13 (4):219-226.
- Alfawaz M. A., 2004 Chemical composition and oil characteristics of pumpkin (*Cucurbita maxima*) seed kernels. Res. Bult., No. (129), Food Science and Agriculture Research Center, King Saud University, Saudi Arabia, pp. 5-18.
- Ali A., Al-Asgah N. A., Al-Ogaily S. M., Ali S., 2000 Effect of dietary lipid source on the growth performance and body composition of *Oreochromis niloticus*. Pakistan Veterinary Journal 20(2):57-63.
- AOAC, 1990 Official methods of analysis of the Association of Official Analytical Chemists. Association of Official Analytical Chemists, 15<sup>th</sup> ed., Arlington, VA, USA, 1298 pp.
- Ayadi F. Y., Rosentrater K. A., Muthukumarappan K., 2012 Alternative protein sources for aquaculture feeds. Journal of Aquaculture Feed Science and Nutrition 4(1):1-26.
- Bahnasawy M. H., 2009 Effect of dietary protein levels on growth performance and body composition of monosex Nile tilapia, *Oreochromis niloticus* L. reared in fertilized tanks. Pakistan Journal of Nutrition 8(5):674-678.
- Bureau of Agricultural Sciences (BAS), 2013 Fish: retail prices by geolocation, commodity, period and year. CountrySTAT Philippines Philippines [online]. [viewed 2014 March 25]. Available from: http://countrystat.bas.gov.ph/.
- Cheng Z. J., Hardy R. W., Usry J. L., 2003 Effects of lysine supplementation in plant proteinbased diets on the performance of rainbow trout (*Oncorhynchus mykiss*) and apparent digestibility coefficients of nutrients. Aquaculture 215:255-265.
- El-Sayed A. F. M., 2006 Tilapia culture. CABI Publishing, Cambridge, MA, 277 pp.
- Food and Agriculture Organization (FAO), 2001 Aquaculture topics and activities: In FAO Fisheries and Aquaculture Department [online]. [viewed 2014 Jan 14]. FAO, Rome, Italy. Available from: http://www.fao.org/fishery/topic/13530/en/.
- Food and Agriculture Organization (FAO), 2006 Contribution to food security and access to food. In State of World Agriculture: FAO Fisheries Technical Paper No. 500, FAO, Rome, pp. 39-52.
- Glew R. H., Glew R. S., Chuang L.-T., Huang Y.-S., Millson M., Constans D., Vanderjagt D. J., 2006 Amino acid, mineral and fatty acid content of pumpkin seeds (*Cucurbita* spp) and *Cyperus esculentus* nuts in the Republic of Niger. Plant Foods for Human Nutrition 61:51-56.
- Haciseferogullari H., Acaroglu M., 2012 Energy balance on pumpkin seed production. Journal of Agricultural Science and Applications 1(2):49-53.
- Hayden B., 1990 Nimrods, piscators, pluckers, and planters. The emergence of food production. Journal of Anthropological Archaeology 9:31-69.
- Huntington T. C., Hasan M. R., 2009 Fish as feed inputs for aquaculture practices, sustainability and implications: a global synthesis. In: Fish as feed inputs for aquaculture: practices, sustainability and implications. Hasan M. R., Halwart M. (eds), FAO Fisheries and Aquaculture Technical Paper No. 518, FAO, Rome, pp. 1-61.
- Kaushik S. J., Coves D., Dutto G., Blanc D., 2004 Almost total replacement of fish meal by plant protein sources in the diet of a marine teleost, the European seabass, *Dicentrarchus labrax.* Aquaculture 230:391-404.
- Khare C. P., (ed) 2007 Indian medicinal plants; an illustrated dictionary. Springer, pp. 182-184, ISBN: 978-0-387-70637-5.
- Kim M. Y., Kim E. J., Kim Y. N., Choi C., Lee B. H., 2012 Comparison of the chemical compositions and nutritive values of various pumpkin (*Cucurbitaceae*) species and parts. Nutrition Research and Practice 6(1):21-27.

- Kirbaslar F. G., Turker G., Ozsoy-Gunes Z., Unal M., Dulger B., Ertas E., Kizilkaya B., 2012 Evaluation of fatty acid composition, antioxidant and antimicrobial activity, mineral composition and calorie values of some nuts and seeds from Turkey. Records of Natural Products 6(4):339-349.
- Magouz F. I., El-Gendi M. O., Salem M. F. I., Elazab A. A., 2008 Use of cucumber, squash and broad bean leaves as non-conventional plant protein sources in Nile tilapia (*Oreochromis niloticus*) diet. 8<sup>th</sup> International Symposium on Tilapia in Aquaculture, pp. 847-859. Retrieved 2013 June 5 from: http://ag.arizona.edu/azagua/ista/ISTA8/ProceedingsISTA8.htm .

Mensah E. T. D., Attipoe F. K., 2013 Growth parameters and economics of tilapia cage culture

- using two commercial fish diets. International Journal of Development and Sustainability 2(2):825-837.
- Radovich T., 2010 Farm and forestry production and marketing for pumpkin and squash (*Cucurbita* spp.). In: Specialty Crops for Pacific Island Agroforestry. Elevitch C. R. (ed), Permanent Agriculture Resouces (PAR), Hawaii, pp. 1-13.
- Raganathan V., Selvasubramanian S., Vasanthakumar S., 2013 Estimation of humoral immune response in rabbits fed with *Cucurbita maxima* seeds. Vet World 6(7): 396-399.
- Rowe D. K., Dean T. L., 1998 Effects of turbidity on the feeding ability of the juvenile stage of six New Zealand freshwater fish species. New Zealand Journal of Marine and Freshwater Research 32:21-29.
- Sadekarpawar S., Parikh P., 2013 Gonadosomatic and hepatosomatic indices of freshwater fish *Oreochromis mossambicus* in response to a plant nutrient. World Journal of Zoology 8(1):110-118.
- Smith G. I., Atherton P., Reeds D. N., Mohammed B. S., Rankin D., Rennie M. J., Mittendorfer B., 2011 Dietary omega-3 fatty acid supplementation increases the rate of muscle protein synthesis in older adults: a randomized control trial. The American Journal for Clinical Nutrition 93:402-412.
- Srbinoska M., Hrabovski N., Rafajlovska V., Sinadinovic-Fiser S., 2012 Characterization of the seed and seed extracts of the pumpkins *Cucurbita maxima* D. and *Cucurbita pepo* L. from Macedonia. Macedonian Journal of Chemistry and Chemical Engineering 31(1):65-78.
- Tacon A. G. J., Metian M., 2008 Global overview on the use of fish meal and fish oil in industrially compounded aquafeeds: trends and future prospects. Aquaculture 285:146-158.
- Yildiz M., 2004 The study of fillet quality and the growth performance of rainbow trout (*Oncorhyncus mykiss*) fed with diets containing different amounts of vitamin E. Turkish Journal of Fisheries and Aquatic Sciences 4:81-86.

Received: 18 March 2014. Accepted: 10 April 2014. Published online: 15 April 2014. Authors:

Marie Noelle M. Fernandez, Department of Biology, School of Science and Engineering, Loyola Schools, Ateneo de Manila University, Katipunan Avenue, Loyola Heights, Quezon City 1108, Metro Manila, Philippines, e-mail: mnm.fernandez@yahoo.com

Janice A. Ragaza, Department of Biology, School of Science and Engineering, Loyola Schools, Ateneo de Manila University, Katipunan Avenue, Loyola Heights, Quezon City 1108, Metro Manila, Philippines, e-mail: jragaza@ateneo.edu

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Teves J. F. C., Fernandez M. N. M., Ragaza J. A., 2014 Effects of replacing fishmeal with squash seed meal (*Cucurbita maxima*) on performance of juvenile Nile tilapia (*Oreochromis niloticus*). AACL Bioflux 7(2):68-75.

Jonni Fay C. Teves, Department of Biology, School of Science and Engineering, Loyola Schools, Ateneo de Manila University, Katipunan Avenue, Loyola Heights, Quezon City 1108, Metro Manila, Philippines, e-mail: jonni\_15\_teves@yahoo.com