

Ateneo de Manila University

Archium Ateneo

Biology Faculty Publications

Biology Department

8-2016

Molecular cloning and expression of chitin deacetylase 1 gene from the gills of *Penaeus monodon* (black tiger shrimp)

Katreena P. Sarmiento
Ateneo de Manila University

Vivian A. Panes
Ateneo de Manila University

Mudjekeewis D. Santos
Ateneo de Manila University

Follow this and additional works at: <https://archium.ateneo.edu/biology-faculty-pubs>



Part of the [Biology Commons](#), and the [Microbiology Commons](#)

Recommended Citation

Katreena P. Sarmiento, Vivian A. Panes, Mudjekeewis D. Santos, Molecular cloning and expression of chitin deacetylase 1 gene from the gills of *Penaeus monodon* (black tiger shrimp), *Fish & Shellfish Immunology*, Volume 55, 2016, Pages 484-489, ISSN 1050-4648, <https://doi.org/10.1016/j.fsi.2016.06.025>.

This Article is brought to you for free and open access by the Biology Department at Archium Ateneo. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of Archium Ateneo. For more information, please contact oadrcw.ls@ateneo.edu.



Short communication

Molecular cloning and expression of chitin deacetylase 1 gene from the gills of *Penaeus monodon* (black tiger shrimp)Katreena P. Sarmiento^{a, b}, Vivian A. Panes^b, Mudjekeewis D. Santos^{a, *}^a Genetic Fingerprinting Laboratory, National Fisheries Research and Development Institute, Mother Ignacia Ave., South Triangle, Quezon City, Metro Manila, 1103, Philippines^b Ateneo de Manila University, Katipunan Ave., Loyola Heights, Quezon City, Metro Manila, 1108, Philippines

ARTICLE INFO

Article history:

Received 8 March 2016

Received in revised form

14 June 2016

Accepted 18 June 2016

Available online 19 June 2016

Keywords:

Chitin deacetylase

Penaeus monodon

White spot syndrome virus

Gills

ABSTRACT

Chitin deacetylases have been identified and studied in several fungi and insects but not in crustaceans. These glycoproteins function in catalyzing the conversion of chitin to chitosan by the hydrolysis of N-acetamido bonds of chitin. Here, for the first time, the full length cDNA of chitin deacetylase (CDA) gene from crustaceans was fully cloned using a partial fragment obtained from a transcriptome database of the gills of black tiger shrimp *Penaeus monodon* that survived White Spot Syndrome Virus (WSSV) infection employing Rapid Amplification of cDNA Ends (RACE) PCR. The shrimp CDA, named *PmCDA1*, was further characterized by *in silico* analysis, and its constitutive expression determined in apparently healthy shrimp through reverse transcription PCR (RT-PCR). Results revealed that the *P. monodon* chitin deacetylase (*PmCDA1*) is 2176 bp-long gene with an open reading frame (ORF) of 1596 bp encoding for 532 amino acids. Phylogenetic analysis revealed that *PmCDA1* belongs to Group I CDAs together with CDA1 and CDA2 proteins found in insects. Moreover, *PmCDA1* is composed of a conserved chitin-binding peritrophin-A domain (CBD), a low-density lipoprotein receptor class A domain (LDL-A) and a catalytic domain that is part of CE4 superfamily, all found in group I CDAs, which are known to serve critical immune function against WSSV. Finally, high expression of *PmCDA1* gene in the gills of apparently healthy *P. monodon* was observed suggesting important basal function of the gene in this tissue. Taken together, this is a first report of the full chitin deacetylase 1 (CDA1) gene in crustaceans particularly in shrimp that exhibits putative immune function against WSSV and is distinctly highly expressed in the gills of shrimp.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

The chitin exoskeleton of crustaceans is an important barrier against the external environment including bacteria and viruses, as well as for maintaining hemolymph and tissue integrity [1]. To form chitin, Chitin deacetylases (CDA) is needed as they aid in the catalysis of the acetamido group in the N-acetyl-D-glucosamine units of chitin [2,3]. CDA is a member of the carbohydrate esterase 4 (CE4) superfamily, sharing the NodB homology domain or polysaccharide deacetylase domain, a conserved domain in the family's primary structure [4]. CDA has been associated with various biological roles. In marine bacteria, CDA function for: chitin catabolism

supporting growth, adaptation to environmental nutrient gradients, tolerance to stress, and protection from predators; and cellular communication [5]. In fungi, CDA facilitate: cell wall synthesis and spore wall formation by increasing chitosan availability, which is a critical component that allows structural rigidity and resistance to various stresses of spores [6]. In insects, CDA are involved in tracheal development and tube elongation [7,8]. To date, however, CDA has never been fully cloned in crustaceans and has never been reported in shrimps.

One of the most lethal viruses that affect shrimp is the white spot syndrome virus (WSSV). It infects most shrimps by traversing into the nuclei of host cells causing disintegration and loss of cellular structure [9]. This virus can cause complete collective mortality within a week that makes a great burden in shrimp agriculture. Because of this, studies that focus on understanding WSSV and host interaction have been carried out including the generation of transcriptome databases that could shed light on

* Corresponding author. National Fisheries Research and Development Institute, Room 601 Corporate 101 Bldg., Mother Ignacia Avenue, Quezon City, 1103, Philippines.

E-mail address: mudjiesantos@gmail.com (M.D. Santos).

genes that are involved in virus pathogenesis [9,10].

Expressed sequence tags (EST) libraries and transcriptome databases serve as a pool of genetic information that may contain valuable information in discovering significant novel genes. More so when there is differential gene expression analysis between healthy and infected organism. In this study, a partial fragment obtained from a transcriptome database extracted from the gills of black tiger shrimp (*Penaeus monodon*) that survived WSSV challenge [10] was fully cloned and identified to be CDA. This study represents the first report on the presence of CDA gene in crustaceans. In addition, it provides the structure and domains of the gene and its putative function, as well as its distinctly high basal expression in gills of shrimp.

2. Materials and methods

2.1. Sample collection

For the cloning part of this study, a live adult *P. monodon* was collected from a local market in Pasay City, Philippines. For constitutive expression analysis in different tissues, three (3) live, apparently healthy *P. monodon* were collected from shrimp farms in Vitali Island, Zamboanga City, Zamboanga Sibugay, and Barangay Sta. Monica, Hagonoy, Bulacan, Philippines. These 3 samples were tested negative for WSSV by Dr. Mary Beth Maningas of the Biochemistry Laboratory, University of Santo Tomas, Philippines using her Juan Amplifier WSSV Diagnostic Kit (JAMP Kit). Upon collection, shrimps were immediately dissected and tissues including: gills, hepatopancreas, intestine, lymphoid organ, heart and hemolymph were collected then preserved in microcentrifuge tube containing RNA later.

2.2. Sequencing of full length *PmCDA1* cDNA gene

Partial sequence was obtained from transcriptome database of *P. monodon* that survived WSSV challenge [10] available at NCBI Sequence Read Archive with accession number SRR57708030 (<http://www.ncbi.nlm.nih.gov/sra>). Total RNA from gills of *P. monodon* was extracted using QIAzol[®] Lysis Reagent (QIAGEN) following the manufacturer's protocol. Subsequently, extracted RNA samples were subjected to RNA purification using RNEasy Mini Cleanup Kit (QIAGEN) following the manufacturer's protocol. RNA yield was determined using Implen[™] nanophotometer.

In the first strand cDNA synthesis, RNA template had a concentration of 0.103 µg/µl. The 10 µl reaction mix had the total concentration as follows: 0.5X first strand buffer, 2 mM DTT, 0.1 mM dNTP mix, SMARTer Oligo IIA (for 5' cDNA synthesis), 5 U RNase inhibitor, 10 U SMARTscribe reverse transcriptase, and 1 mM of respective primers for 5' and 3' cDNA synthesis. The reaction mix was subjected to 42 °C for 90 min and heated to 70 °C for 10 min in a thermal cycler. Subsequently, the samples were diluted with 20 µl of tricine-EDTA buffer.

On the other hand, RACE was carried out with the in a 25 µl reaction mix with final concentration as follows: 1X PCR buffer, 1 mM dNTP mix, 1X polymerase mix, 1X universal primer, and gene specific primers. Specific primers designed for facilitating 5' (*PmCDA1* 5RACE-1 and *PmCDA1* 5RACE-2) and 3' RACE (*PmCDA1* 3RACE-1) extensions are shown in Table 1. Primer annealing sites are shown in Fig. S1. The reaction mixes were subjected to the following conditions: initial denaturation at 94 °C for 3 min, 38 cycles of 94 °C for 30 s, 68 °C for 30 s and 72 °C for 3 min, then final extension at 72 °C for 10 min. All PCR amplicons were visualized on a 1% gel stained with ethidium bromide and documented with SynGene G:BOX. Positive amplicons were sent to Macrogen Inc. (Korea) for sequencing.

Table 1

Primer sequences used to amplify and sequence chitin deacetylase 1 in *P. monodon* cDNA.

Primer name	Sequence (5' → 3')
<i>PmCDA1</i> 5RACE-1	TTCTTCCACTCTGCCTTCCTG
<i>PmCDA1</i> 5RACE-2	TAGCAGTAGCAATCGGGGAGGGCGCAG
<i>PmCDA1</i> 3RACE-1	GCGTCGTCGACGGCCACCCCTTC
GSP forward	GTCATGAACGAGATGGACCG
GSP reverse	CAGTACTTGCTCTACCTGGCC
EF1 α forward	ATGGTTGTCAACTTTGCCCC
EF1 α reverse	TTGACCTCTTGATCACACC

2.3. In silico analysis

The nucleotide sequence, translated amino acids, and average molecular weight were analyzed and determined using Geneious 6.1.8 (Biomatters). SignalP 4.1 (<http://www.cbs.dtu.dk/services/SignalP/>) and NetNGlyc 1.0 (<http://www.cbs.dtu.dk/services/NetNGlyc/>) were utilized to predict signal peptide cleavage and N-glycosylation sites, respectively. Nucleotide and amino acid sequence identities were determined using BLASTn and BLASTp, respectively. Complete multiple alignments were conducted using ClustalW with default parameters. Conserved Domain Architecture Retrieval Tool (CDART) (<http://www.ncbi.nlm.nih.gov/Structure/lexington/lexington.cgi>) was used to determine conserved domains. Sequences of the members of CE4 superfamily (GENBANK: cl15692) were retrieved from NCBI. Predicted 3-dimensional structure of the protein was generated through Protein Homology/analogy Recognition Engine V2.0 (Phyre²) (www.sbg.bio.ic.ac.uk/phyre2/). Phylogenetic analysis was carried out in MEGA 6 (<http://www.megasoftware.net/>) for the generation of tree using Maximum Likelihood (ML) method based on the Whelan and Goldman (WAG) model [29] with 500 bootstrap tests and complete deletion of sites.

2.4. Expression analysis

Total RNA from different tissues of three (3) apparently healthy *P. monodon* samples were extracted using QIAzol[®] Lysis Reagent (QIAGEN) following the manufacturer's protocol. These are then used for cDNA synthesis. The reverse transcription (RT) reaction mix had the components with final concentration as follows: 1X Buffer RT, 0.5 mM dNTPs, 1 µM Oligo-dT primer, 10 units RNase inhibitor, 4 units Omniscript Reverse Transcriptase (QIAGEN), with 0.203 µg/µl of RNA template. RT – Polymerase Chain Reaction was used to determine expression of *PmCDA1* from different tissues of healthy *P. monodon* following [12] with modifications. Primers used for the amplification of *PmCDA1* (GSP forward and GSP reverse) and elongation factor 1 alpha (EF1 α) (EF1 α forward and EF1 α reverse) are shown in Table 1. The following cycling parameters: initial denaturation at 95 °C for 5 min, 30 cycles of denaturation at 95 °C for 30 s, annealing at 55 °C for 30 s, 72 °C for 30 s, and final extension at 72 °C for 10 min, were used for amplification of both *PmCDA1* and EF1 α . Negative control contained no template. All PCR amplicons were visualized on a 1% gel stained with ethidium bromide and documented with SynGene G:BOX. Relative expression based on light intensity was measured using ImageJ software (<http://imagej.nih.gov/ij/>) [11] and standard deviations were computed.

3. Results and discussion

This study serves as the first report on the presence of chitin deacetylase (CDA) gene in crustaceans particularly in shrimps.

Results revealed that the shrimp CDA (GENBANK: KU156741.1), has a length of 2176 bp with open reading frame (ORF) of 1596 bp encoding for 532 amino acids (AA), and a molecular weight of 61.271 kDa (GENBANK: ALO20448.1) (Fig. 1). BLAST analysis showed 63% identity of protein query with rice grasshopper *Oxya chinensis* chitin deacetylase 1 (*OcCDA1*) and red flour beetle *Tribolium castaneum* chitin deacetylase precursor [13,14] and 59% identity with cotton bollworm *Helicoverpa armigera* chitin deacetylase 1 (*HaCDA1*) [15]. The nucleotide sequence contains 59 bp upstream-untranslated region and 518 bp downstream-untranslated region, followed by poly-adenylation signal sequence (AATAAA) and A + T rich region. Further analysis revealed that it includes a potential signal peptide in its AA positions 23 and 24, and contains N-glycosylation sites at AA positions 236, 260 and 288. Chitin-binding peritrophin-A domain (CBD) (AA41 to AA95), low-density lipoprotein receptor class A domain (LDL-A) (AA 115 to AA 149) and catalytic NodB homology domain of the carbohydrate esterase 4 superfamily (CE4) (CDD accession: cl15692) (AA 191 to AA 460) that includes polysaccharide deacetylase (AA 192 to 301) were also

found. CE4 domain was described to have several signature motifs (TFDD, H[S/T]xxHP, RxP[Y/F], DxxDW, GxxxFxx) that render different functions, and are variable that allows distinction in enzyme activity [16]. The predicted 3D structure of *PmCDA1* is shown in Fig. S2. The model revealed alpha helices and beta strands in the protein structures as well as conserved sites based on Conserved Domain Database (CDD) including putative catalytic sites and NodB motif.

Chitin-binding peritrophin-A domain (CBD), the first conserved domain found, is known to be significant in enhancing enzyme activity to allow prolonged interaction with the substrate [17,18]. revealed that a group of CBD containing proteins is one of the four groups of proteins that are highly up-regulated after WSSV infection suggesting its immune-related function against the virus. CBD-containing proteins were included in the fifty most abundant genes and unique genes with increased differential abundance in WSSV-infected library [9]. This supports the transcriptome database of *P. monodon* that survived WSSV challenge generated by Ref. [10] wherein CBD-containing CDA was differentially expressed in

AAGCAGTGGTATCACGCAGAGTACATGGGGTCAGTGAACGGTCTGGGCTGAGCGTTAAC	59
ATG GCAAGAGTAAGA TGC GG T AGT TCT CTT GCC CTT CTT GGC ATT GTG CTG CCG AGC GGG TCA AG AGG CAG G CAG	134
M A R V R C G S S L A L L G I V L P S G S R G R Q	25
TTT CAGACA CGA CCG AGG ATG GAG CGA ACC TTC AAG AGG GAA CTG TGC AAG GAT AAG GGC GCC GGC GAG TGG TTC	209
F Q T R P R M E R T F K R E L C K D K G A G E W F	50
CGT CTC AGC CTC GGT GAC TGT CGC GAT GTG ATC CAG TGT ACC GAC GCC GGT CTT CAG GCG CTG CGT TGC CCC CAC	284
R L S L G D C R D V I Q C T D A G L Q A L R C P H	75
GGC CTG GCC TTC AAC CTG GAG CAG ACC CTT GCG AC TGG AAA GCC AAC GTC AAG AAC TGC GAC AGG AAG GAG AAG	359
G L A F N L E Q Q T C D W K A N V K N C D R K E K	100
ACG AAG GTC G TGA AG CCT CTC TTC AAC ACC GT C GAG CCC CTT TGC CAG GAA AAC CAG TTG GCG TGC GGT GAC GGC	434
T K V V K P L F N T V E P L C Q E N Q L A C G D G	125
ACC TGT CTG GAT AGG CAG GTC TTC TGT GAT G G A A A G A A G AT TGT ACC GAT GGT TCT GAC GAG ACC GAT GT GAC	509
T C L D R Q V F C D G K E D C T D G S D E T A C D	150
GTG AAAA T GAC CCC AAC AGC GCT CCC ATC T G C A A C C C G A G A C T G C C G C T C C C C G A T T G C T A C T G T A C A A C	584
V K N D P N S A P I C N T E D C R L P D C Y C Y N	175
GAC CCT AGC GAG ATC CCC CACA ACAT GAAG CCT CCG AAG TG CCT CAG ATGG T CACC ATCACT TTC GAC GAT GCC	659
D P S E M P H N M K P S E V P Q M V T I T F D D A	200
ATC AAC ATCAACA ATG GAC CTG TAC GAG CTC ATC TTC AAG CAG CGC TTC AAC CCC AAC GGCT GCT CCA TC AAG	734
I N I N N M D L Y E L I F K Q R F N P N G C S I K	225
TCC ACT TCT TCG TT TCT CAC AAA TAC AAC AACTAC ACC GCC ACT CAG GAG ATG CAC CGT CTG GGT CAC GAG ATC	809
S T F F V S H K Y N N Y T A T Q E M H R L G H E I	250
GCT GTT CACT CCA CAC CAC GCA AAC AAC GAG ACT TTT CTT GTC CAC GCA AGC GAA GAC GAG TAC GAG CGC GAA	884
A V H S I T H A N N E T F W S H A S E D E Y E R E	275
ATG GGT GGT GCC CGC GTC ATC ATC GAA CGC TTC GCC AAC ATC ACC GAC CAG TCC ATC ATC GGC ATG CGT AAT CCC	959
M G G A R V I I E R F A N I T D Q S I I G M R N P	300
TTC CTC CGT GTG GGC CGS AAC AGC CAG TTC CAG GAT GAT G G A A G A A C A C C T T C C T G T A C G A C T C C A C C A T C	1034
F L R V G G N S Q F R M M E K N T F L Y D S T I T	325
GCC CCC CTG TCC TCC ATG CCC CTG TGG CCC TAC ACC CTT GACT ACC CGC ATG CCC CAC CCC TGC CAT G G A A A C CTC	1109
A P L S S M P L W P Y T L Y Y R M P H P C H G N L	350
CAG AAC TG CCCC ACT CGCT CCT TCC GCA GTG TGG GAG ATG GTC ATG AAC GAG ATG GAC CGT CGC GAG GAA ACC ACT	1184
Q N C P T R S F A V W E M V M N E M D R R E E P T	375
TAC GAG GAC GGC CTC CCC GGA TGC CAC ATG GTC GAT TCT TGT TCG CCA CCA AG CCC GAG CCC GAG CAG TTC TAC	1259
Y E D G L P G C H M V D S C F A T K P E P E Q F Y	400
AAC TTC CTG CAG AAC AAC TTC AAC CGC CAC TAC AAG TCA AAC CGC GCT CCC TTC GGT CTG TTC TTC CACT CT GCC	1334
N F L Q N N F N R H Y K S N R A P F G L F F H S A	425
TTC CTG AAG AAC AAC CCC GAC ATC CTG GAC ACC TTC CTCTACT GGT GAC GAG ACC CTG AAG AAC CAGAAG GAC	1409
F L K N N P D I L D T F L Y W L D E T L K N Q K D	450
GTT TACT TCG TACC ATG ACC CAG GTC ATC CAG TGG ATG CAG GAC CCC CGC CCC GTC GGC CAG CTG AACAAC TAC	1484
V Y F V T M T Q V I Q W M Q D P R P V G Q L N Y Y	475
GAG GCCTGGAAGGAGAAGTGCCTGTCGACGGGCCCA CCTTCTGCTACGGCGCAACACTGCGAGCTGGAC ACT	1559
E A W K E K C V V D G P P F C Y G G N N C E L D T	500
GAC GAG CTC CCT GGC CAG ACC CTC CAC CTG TCC ACC TGC ATG CCG TGC CCC AAC AAT TAT CCTTGGACAAG GAC	1634
D E L P G Q T L H L S T C M R C P N N Y P W T R D	525
CCA TTGGCGAGGGAATTCCTTAA AGAGCAGTCGCTGCCGCTTACGCTGCGCCTTGC CCCCTGGGTGAGCCAGGCC	1712
P L G E G F F *	532
CCCTCCCCTAGGGTGCAGTTGCTCACTTGTAAATATTGTTTTATAATTTGTACATCGGCCGCGACCAGCTTGTGTCT	1791
AGGCAAGAGTCGCGGTGTTCCATCTTGACACCCGTCGTGACGTTACGGCGCCATGGCATACCGAGACAGGTGACGCTAGT	1870
TGCCACGGCGAAGAACCATGACAGGCTGGGGCGGGTCTTCCAGTGGGACCAAGCAGGGTTCGAATCCCCGCAAGAAAGAC	1949
CCTAAACCTGGGAAGATTCGAGTGGGGCGGCCCTACCATATTTATTGGAGCTGGAGGACCCCTTTTCCCTCCTTATC	2028
ACACACCCCATAACTTTCATATCCCCCTCCCCCTAGATCTCTCGTCTATATATTTTTTTCTTTCTTTTCTTTTCTTTT	2107
TTTTTTTTTAAATAAA CATATTAATTTTTTTTTTAAACGGGGCCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	2176

Fig. 1. The full-length cDNA sequence and deduced amino acid sequences of *PmCDA1*. The predicted signal peptide is underlined with a broken line. Chitin-binding peritrophin-A domain is highlighted in light gray, low-density lipoprotein receptor class A domain is highlighted in dark gray, and catalytic NodB homology domain of the carbohydrate esterase 4 superfamily is underlined with two lines and signature motifs are boxed with solid line. Predicted N-glycosylation sites are boxed with broken line and poly-adenylation signal is italicized.

survivor shrimp relative to apparently healthy shrimp. CBD is present in insects and animal chitinases as part of peritrophic matrix proteins, including peritrophins [19]. Peritrophin in fleshy prawn *Fenneropenaeus chinensis* was reported to have gram-negative bacteria binding activity and chitin binding activity, and was found to be highly induced in hemocytes, heart, stomach, gut, and gills of infected shrimp, predicting its role in immune defense and other physiological responses [20]. Previously reported *P. monodon* chitin-binding protein (PmCBP) and peritrophin-like protein from pacific white shrimp *Litopenaeus vannamei* (LvPT) revealed interaction between host shrimp and envelope protein of WSSV, which further strengthens the significance of the role of proteins with chitin-binding activity in shrimp-WSSV interaction [21,22]. Collectively, the presence of CBD in the shrimp CDA may also be involved in defense mechanism of *P. monodon* against WSSV.

Low-density lipoprotein receptor class A domain (LDL-A) was the second domain found in the shrimp CDA. In kuruma shrimp *Marsupenaeus japonicus*, C-type lectins (CTLs) which binds to WSSV causing decreased WSSV replication contain critical LDL-A to carry out pattern-recognition mechanisms during immune response [23]. In fact, disrupting CTLs caused increased WSSV replication and allows attachment and penetration of WSSV in shrimp [23]. Given this, LDL-A in shrimp CDA may also be predicted to have the same binding activity of pattern-recognition mechanisms in black tiger shrimp. LDL-A also occurs in insect and fungi CDAs, however no further function of the domain have been discussed in previous reports.

A putative catalytic domain of CDA proteins which serves as part of CE4 superfamily was also observed in shrimp CDA. The CE4 superfamily is comprised of different enzymes including chitin deacetylase, bacterial peptidoglycan *N*-acetylglucosamine deacetylases and acetylxyylan esterases that aids in the catalysis of *N*- or *O*-deacetylation of their respective substrates [24]. Among the AA sequence of these enzymes, the NodB homology domain is always present. Previous reports about the domain mainly focuses on nitrogen-fixing bacteria that mediates synthesis of host-specific signal molecules [25]. However, the function of this domain has not been extensively studied in other organisms.

CDAs are suggested to occur from five to nine members, categorized into five groups (Group I – V), depending on the species [30]. These CDAs have variable assortment of domains that sometimes differ in organization. Group I includes CDA1 and CDA2 that have single and multiple isoforms, respectively. CDAs belonging to this group have a complete set of the three domains: CBD, LDL-A, and CE4. Group II includes CDA3 that has similar domain organization with CDAs from group I but are substantively different in the amino acid composition. Group III includes CDA4 that lacks LDL-A. Group IV includes CDA5 with similar organization with CDA4 only that CBD and CE4 are separated by a long glutamine-rich intervening region. Group V includes CDA5–9 that lacks CBD and LDL-A, then are only left with CE4. Since the shrimp CDA possess a complete set of CBD, LDL-A, and CE4 domains (Fig. S3), it can be grouped to Group 1.

Phylogenetic analysis using ML method involving eighteen (18) AA sequences with a total of 335 positions also revealed that the shrimp CDA belonged to group I CDAs, forming a clade together with reference sequences of CDA1 and CDA2 from various insects supported by a bootstrap value of 100% (Fig. 2). Previous studies revealed that CDA1 are found in all insect species whereas CDA2 are specific only to pupa, and that pre-mRNAs of CDA2 undergo RNA splicing and/or exon skipping to yield CDA isoforms [30,31]. These two CDAs were also distinctly differentiated based on their signature motifs in its CE4 domain, particularly TFDD motif 1 in CDA1 was substituted as TFNG in all isoforms of CDA2 that makes it




Fig. 2. Phylogenetic analysis of *PmCDA1* and amino acid sequences of CE4 super family CDA members using ML method, and WAG model. The percentage of trees in which the associated taxa clustered together is shown next to the branches. Reference CDA sequences were obtained from the following organisms: red flour beetle *Tribolium castaneum* (Tc), cotton bollworm *Helicoverpa armigera* (Ha), brown planthopper *Nilaparvata lugens* (NI), rice grasshopper *Oxya chinensis* (Oc), and silkworm *Bombyx mori* (Bm). All reference sequences obtained from GENBANK were labelled with database accession numbers.

inactive [30]. CDA from *P. monodon* was found to have TFDD motif 1 (Fig. 1) suggesting that it is CDA1 hence it is named here as *PmCDA1*. Differences in motifs from CDAs of different groups are further exemplified in multiple alignment of CDA sequences (Fig. S4).

The present study also observed that *PmCDA1* was specifically highly expressed in the gills of black tiger shrimp (Fig. 3). Gills in shrimps serve as the predominant site for the formation of hemocyte nodules during injection of foreign particles and accumulation of viable bacteria during infection, suggesting its significant role in shrimp defense [32,33]. Gills mediate direct contact to the external environment, hence its cells play a vital role in response to biotic and abiotic factors [34]. In addition to this, *PmCDA1* was also observed to be expressed in the muscle and lymphoid organ of shrimps. The lymphoid organ in penaeid prawns was predicted to be a major phagocytic organ that serves a major site for viral degradation [26,27]. Previous study revealed that muscle-related genes were upregulated after pathogen infection which serves as an effective way in clearing invading pathogens [28].

Expression of CDA from other species was reported to be high during early developmental stages of insects and pathogen infection in fungi. In rice grasshopper *O. chinensis*, highest expression of *OcCDA1* was observed in the integument of early stage of its development [13]. In addition, expression of *NICDA1* peaked in the moulting stage of the brown planthopper *Nilaparvata lugens* suggesting its critical role in the turnover of the old insect cuticle [31]. Upregulation of CDA gene in the common mushroom *Agaricus bisporus* infected with mycoparasite *Verticillium fungicola* was previously observed [35]. In contrast, down-regulation of chitin deacetylase-like protein was observed during baculovirus infectivity in cotton bollworm *H. armigera*, which serves as a defense mechanism preventing entrance of baculoviruses in its host by decreasing peritrophic membrane permeability [15]. Given this, and the distinct high expression of *PmCDA1* in gills of apparently healthy shrimp suggest an important functional role for the said gene at a possible entry site of infection. Such is worth pursuing in future studies to elucidate further “resistance” of WSSV “survivor”




Fig. 3. 1. Gel electrophoresis of constitutive expression analysis of black tiger shrimp *PmCDA1* in various tissues relative to expression of EF1- α as determined by RT-PCR. 2. Constitutive expression of black tiger shrimp *PmCDA1* in various tissues relative to expression of EF1- α as determined by RT-PCR and ImageJ. The bars represent mean values of three shrimp samples plus Standard Deviation (SD).

shrimps.

In conclusion, the study fully cloned and identified the first CDA in crustaceans, particularly in black tiger shrimp *P. monodon*, named here as *PmCDA1*. In addition, it provided the structure and domains of the gene and its possible role in the immune function of the gills of shrimp against WSSV. Such discovery is important in further understanding the effects of WSSV in the immune system of black tiger shrimp, an economically important culture species worldwide, and in providing long-term molecular-based solutions in combating the WSSV infection.

Authors' contributions

KS carried out the molecular genetic studies, participated in the sequence alignment, and drafted the manuscript. VP helped in drafting the manuscript. MS conceived the study, participated in its design and coordination, and structured the writing of the manuscript. All authors read and approved the final manuscript.

Conflict of interests

The authors declare that they have no competing interests.

Acknowledgements

The authors would like to thank Mr. Muhaimin Albani of the City Agriculturist of Zamboanga for facilitating the sampling of shrimps from Vitali, Zamboanga City and Dr. Mary Beth Maningas and colleagues from the University of Santo Tomas, Manila for testing the said shrimp samples for WSSV infection. We are also thankful to Ms. Jacqueline Marjorie R Pereda, Ms. Minerva Fatimae H Ventolero

and all the members of GFL for the support. Finally, the authors would also like to thank the National Fisheries Research and Development Institute (NFRDI) for its funding and laboratory support. We are also thankful to the Department of Science and Technology – Accelerated Science and Technology Human Resource and Development Program as well as their Student Research Support Fund for supporting the accomplishment of the project.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.fsi.2016.06.025>.

References

- [1] A. Tassanakajon, K. Somboonwiwat, P. Supungul, S. Tang, Discovery of immune molecules and their crucial functions in shrimp immunity, *Fish Shellfish Immunol.* (2012) 1–14, <http://dx.doi.org/10.1016/j.fsi.2012.09.021>.
- [2] E. Cohen, Chitin biochemistry: synthesis, hydrolysis, and inhibition, *Adv. Insect Physiol.* 38 (2010) 5–74.
- [3] Y. Araki, E. Ito, A pathway of chitosan formation in *Mucor rouxii*, *Eur. J. Biochem.* 55 (1975) 71–78.
- [4] P.M. Coutinho, B. Henrissat, Carbohydrate-active enzymes: an integrated database approach, in: H.J. Gilbert, G. Davies, B. Henrissat, B. Svensson (Eds.), *Recent Advances in Carbohydrate Bioengineering*, The Royal Society of Chemistry, Cambridge, 1999, pp. 3–12.
- [5] Y. Zhao, R.D. Park, R.A.A. Muzzarelli, Chitin deacetylases: properties and applications, *Mar. Drugs* 8 (2010) 24–46.
- [6] M.E. Gilmore, D. Bandyopadhyay, A.M. Dean, S.D. Linnstaedt, D.L. Popham, Production of muramic delta-lactam in *Bacillus subtilis* spore peptidoglycan, *J. Bacteriol.* 186 (2004) 80–89.
- [7] S. Luschign, T. Bätz, K. Armbruster, M.A. Krasnow, Serpentine and vermiform encode matrix proteins with chitin binding and deacetylation domains that limit tracheal tube length in *Drosophila*, *Curr. Biol.* 16 (2006) 186–194.
- [8] S. Wang, S.A. Jayaram, J. Hemphälä, K.A. Senti, V. Tsarouhas, H. Jin, C. Samakovlis, Septate-junction-dependent luminal deposition of chitin deacetylases restricts tube elongation in the *Drosophila* trachea, *CurrBiol* 16 (2006) 180–185.
- [9] W. Leu, Y. Wang, D. Tian, Z. Yin, J. Kwang, Detection of white spot syndrome virus (WSSV) of shrimp by means of monoclonal antibodies (MAbs) specific to an envelope protein (28 kDa), *Dis. Aquat. Org.* (2002) 11–18.
- [10] B. Maralit, M. Ventolero, M. Maningas, E. Amar, M. Santos, Subtracted transcriptome profile of black tiger shrimp (*Penaeus monodon*) that survived WSSV challenge, *Dataset Pap. Sci.* 2014 (2014) 1–11.
- [11] M.D. Abramoff, P.J. Magalhaes, S.J. Ram, Image processing with ImageJ, *Biophot. Int.* 11 (2004) 36–42.
- [12] M. Santos, M. Yasuike, I. Hirono, T. Aoki, The granulocyte colony-stimulating factors (CSF3s) of fish and chicken, *Immunogenetics* (2006) 422–432.
- [13] G.W. Ding, R.R. Yu, M.L. Yang, E.B. Ma, J. Yang, Zhang, Molecular characterization and function analysis of chitin deacetylase 1 gene in *Oxya chinensis* (Orthoptera: acrididae), *Acta Entomol. Sin.* 57 (11) (2014) 1265–1271.
- [14] S. Jaspurua, Y. Arakane, G. Osman, K.J. Kramer, R.W. Beeman, S. Muthukrishnan, Genes encoding proteins with peritrophin A-type chitin-binding domains in *Tribolium castaneum* are grouped into three distinct families based on phylogeny, expression and function, *Insect Biochem. Mol. Biol.* 40 (3) (2010) 214–227.
- [15] A.K. Jakubowska, S. Caccia, K.H. Gordon, J. Ferre, S. Herrero, Downregulation of a chitin deacetylase-like protein in response to baculovirus infection and its application for improving baculovirus infectivity, *J. Virol.* 84 (5) (2010) 2547–2555.
- [16] D.E. Blair, A.W. Schuttelkopf, J.I. MacRae, D.M. Van Aalten, Structure and metal-dependent mechanism of peptidoglycan deacetylase, a streptococcal virulence factor, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 15429–15434.
- [17] B.L. Cantarel, P.M. Coutinho, C. Rancurel, T. Bernard, V. Lombard, B. Henrissat, The Carbohydrate-Active EnZymes database (CAZy): an expert resource for glycogenomics, *Nucleic Acids Res.* 37 (2009) D233–D238.
- [18] J.H. Leu, S.H. Chen, Y.B. Wang, Y.C. Chen, S.Y. Su, C.Y. Lin, J.M. Ho, C.F. Lo, A review of the major penaeid shrimp EST studies and the construction of a shrimp transcriptome database based on the ESTs from four penaeid shrimp, *Mar. Biotechnol.* (2010), <http://dx.doi.org/10.1007/s10126-010-9286-y>.
- [19] Z. Shen, M. Jacobs-Lorena, A type I peritrophic matrix protein from the malaria vector *Anopheles gambiae* binds to chitin. Cloning, expression, and characterization, *J. Biol. Chem.* 273 (1998) 17665–17670.
- [20] X.J. Du, J.X. Wang, N. Liu, X.F. Zhao, F.H. Li, J.H. Xiang, Identification and molecular characterization of a peritrophin-like protein from fleshy prawn (*Fenneropenaeus chinensis*), *Mol. Immunol.* 43 (2006) 1633–1644, <http://dx.doi.org/10.1016/j.molimm.2005.09.018>.
- [21] S. Xie, X. Zhang, J. Zhang, F. Li, J. Xiang, Envelope proteins of white spot syndrome virus (WSSV) interact with *Litopenaeus vannamei* peritrophin-like

- protein (LvPT), PLoS One 10 (12) (2015) e0144922, <http://dx.doi.org/10.1371/journal.pone.0144922>.
- [22] K.Y. Chen, T.C. Hsua, P.Y. Huang, S.T. Kang, C.F. Loc, W.P. Huang, L.L. Chen, *Penaeus monodon* chitin-binding protein (PmCBP) is involved in white spot syndrome virus (WSSV) infection 27 (3) (2009) 460–465.
- [23] Y.H. Xu, W.J. Bi, X.W. Wang, Y.R. Zhao, X.F. Zhao, J.X. Wang, Two novel C-type lectins with a low-density lipoprotein receptor class A domain have antiviral function in the shrimp *Marsupenaeus japonicus*, Dev. Comp. Immunol. 42 (2) (2014) 323–332, <http://dx.doi.org/10.1016/j.dci.2013.10.003>, Epub 2013 Oct 16.
- [24] A. Marchler-Bauer, M.K. Derbyshire, N.R. Gonzales, S. Lu, F. Chitsaz, L.Y. Geer, R.C. Geer, J. He, M. Gwadz, D.I. Hurwitz, C.J. Lanczycki, F. Lu, G.H. Marchler, J.S. Song, N. Thanki, Z. Wang, R.A. Yamashita, D. Zhang, C. Zheng, S.H. Bryant, CDD: NCBI's conserved domain database, Nucleic Acids Res. 43 (Database issue) (2015) D222–D226, <http://dx.doi.org/10.1093/nar/gku1221>, Epub 2014 Nov 20.
- [25] M. John, H. Röhrig, J. Schmidt, U. Wieneke, J. Schell, Rhizobium NodB protein involved in nodulation signal synthesis is a chitoooligosaccharide deacetylase, Proc. Natl. Acad. Sci. U. S. A. 90 (1993) 625–629, <http://dx.doi.org/10.1073/pnas.90.2.625>, Biochemistry.
- [26] M.S. Anggraeni, L. Owens, The haemocytic origin of lymphoid organ spheroid cells in the penaeid prawn *Penaeus monodon*, Dis. Aquat. Org. 40 (2000) 85–92.
- [27] M. Kondo, T. Itami, Y. Takahashi, R. Fujii, S. Tomonaga, Structure and function of the lymphoid organ in the kuruma prawn, Dev. Comp. Immunol. 18 (Suppl. 1) (1994) S109.
- [28] S. Nayak, S.K. Singh, N. Ramaiah, R.A. Sreepada, Identification of upregulated immune-related genes in *Vibrio harveyi* challenged *Penaeus monodon* post-larvae, Fish Shellfish Immunol. 29 (3) (2010) 544–549.
- [29] S. Whelan, N. Goldman, A general empirical model of protein evolution derived from multiple protein families using a maximum-likelihood approach, Mol. Biol. Evol. 18 (5) (2001) 691–699.
- [30] R. Dixit, Y. Arakane, C.A. Specht, C. Richard, K.J. Kramer, R.W. Beeman, S. Muthukrishnan, Domain organization and phylogenetic analysis of proteins from the chitin deacetylase gene family of *Tribolium castaneum* and three other species of insects, Insect. Biochem. Mol. Biol. 38 (2008) 440–451.
- [31] Y. Xi, P.L. Pan, Y.X. Ye, B. Yu, C.X. Zhang, Chitin deacetylase family genes in the brown planthopper, *Nilaparvata lugens* (Hemiptera: Delphacidae), Insect. Mol. Biol. 23 (2014) 695–705.
- [32] G. Martin, M. Quintero, M. Quigley, H. Khosrovian, Elimination of sequestered material from the gills of decapod crustaceans, J. Crustac. Biol. 20 (2000) 209–217.
- [33] Burgents JE, Burnett LE, Stabb EV, Burnett KG. Localization and bacteriostasis of *Vibrio* introduced into the Pacific white shrimp, *Litopenaeus vannamei*. Dev. Compar. Immunol. 2.
- [34] W. Wu, B. Wu, T. Ye, et al., TCTP is a critical factor in shrimp immune response to virus infection, in: J. Spencer (Ed.), PLoS ONE 8(9), 2013 e74460, <http://dx.doi.org/10.1371/journal.pone.0074460>.
- [35] P. Mills, J. Thomas, M. Sergeant, A. Costa, P. Collopy, A. Bailey, G. Foster, M. Challen, Chapter 1 Interactions between *Agaricus bisporus* and the pathogen *Verticillium fungicola*, Br. Mycol. Soc. Symp. Ser. (2008) 1–17.